

Dipole and Bloch oscillations of cold atoms in a parabolic lattice

A. V. Ponomarev and A. R. Kolovsky

Kirensky Institute of Physics, Ru-660036 Krasnoyarsk, Russia

(Dated: February 2, 2008)

The paper studies the dynamics of a Bose-Einstein condensate loaded into a 1D parabolic optical lattice, and excited by a sudden shift of the lattice center. Depending on the magnitude of the initial shift, the condensate undergoes either dipole or Bloch oscillations. The effects of dephasing and of atom-atom interactions on these oscillations are discussed.

1. Bloch oscillations (BO) of a quantum particle in a periodic potential are one of the most fascinating phenomena of quantum physics [1]. Since the pioneering experiment [2] in 1996, this phenomenon has been intensively studied for cold atoms in optical lattices [3], with recent emphasis on quantum statistical (Fermi or Bose) and atom-atom interaction effects. In particular, the dynamics of degenerate Bose gases, on which we will focus here, was studied experimentally in [4, 5, 6]. It should be stressed from the very beginning that, when addressing this problem theoretically, one has to distinguish between quasi one-dimensional lattices (created by two counter-propagating laser beams) and truly 1D lattices (or so-called modulated quantum tubes). Indeed, in the former case the number of atoms per well of the optical lattice can be as large as $10^3 - 10^4$, and a mean field approach (based on the Gross-Pitaevskii or nonlinear Schrödinger equation) is generally justified. This is not the case of the truly 1D lattices, where only few atoms occupy a single well, and, hence, a microscopic analysis is required. For a tilted infinite lattice such analysis, based on the Bose-Hubbard model, was presented in [7, 8, 9], where two regimes of BO – quasiperiodic and irreversible decaying – were identified.

When referring to the typical laboratory experiments, an additional complication stems from the harmonic confinement along the lattice. Clearly, harmonic confinement should modify BO of bosonic atoms, and the aim of this work is to estimate its effect. At the same time, parabolic lattices have their own interest, because they allow to study dipole oscillations of BECs. Recent experiments [10] have shown that there is a fundamental difference between dipole oscillations in quasi- and truly 1D lattices. While in the former case the main effect of the periodic potential can be taken into account by simply substituting the atomic mass by its effective mass in the ground Bloch band [11], one observes a rapid decay of oscillations in the latter case. In the present paper we also briefly discuss dipole oscillations of a BEC in truly 1D lattices, partially overlapping in this part with recent theoretical work [12].

2. We consider atoms in a parabolic lattice potential $V(x) = M\omega^2 x^2/2 - V \cos^2(2\pi x/d)$, where the atoms are set into motion by a sudden shift of the trap origin. The relevant parameters of the system are the hopping matrix element J (defined by the amplitude of the periodic potential V), the ‘parabolicity’ $\nu = M\omega^2 d^2$, and the initial shift $l_0 = \Delta x/d$. Using the single band approximation

and neglecting atom-atom interactions, the dynamics of the system is described by the pendulum model [3, 13],

$$i\hbar\dot{a}_l = \frac{\nu}{2}l^2 a_l - \frac{J}{2}(a_{l+1} + a_{l-1}), \quad (1)$$

where $a_l(t)$ is the complex amplitude of the atoms in the l -th well of the optical lattice. The separatrix of the pendulum corresponds to the shift

$$l^* = 2(J/\nu)^{1/2}. \quad (2)$$

If the initial shift $l_0 < l^*$, the pendulum shows oscillations around the equilibrium point and, referring to the original system, this regime is regarded as that of dipole oscillations of the atoms. If $l_0 > l^*$, the pendulum is in the rotational regime, and the dynamics of the atoms can be regarded as BO in a local static field $F = \nu l_0/d$.

3. We begin with analyzing the regime of BO for an ideal bosonic system (i.e., no atom-atom interactions). Because the local static force F is not homogeneous, one has an additional process of dephasing of BO in a parabolic lattice, as compared to the paradigmatic case of a homogeneously tilted lattice. When discussing the mean atomic momentum, we can estimate this effect of dephasing by evaluating the sum

$$p(t) \sim \sum_m \exp(-m^2/2\gamma^2) \sin[(\omega_B + \nu m/\hbar)t], \quad (3)$$

where $\omega_B = \nu l_0/\hbar$, $m = l - l_0$, and γ is the width of the atomic wave packet (measured in units of the lattice period). Substituting the sum by an integral, we obtain

$$p(t) \sim \exp(-t^2/2\tau_\gamma^2) \sin(\omega_B t), \quad (4)$$

where

$$\tau_\gamma = \hbar/\gamma\nu. \quad (5)$$

It is seen in Eq. (5), that the dephasing time τ_γ is defined by both the wave packet width and by the trap frequency (see Fig. 1 below). On the basis of this result one might conclude that a narrow wave packet is preferable for studying BO in parabolic lattices. This is, however, not exactly true because a narrow wave packet implies a lower contrast of the interference pattern measured in laboratory experiments. Thus, one has to keep a compromise between the contrast and dephasing, when preparing the initial wave packet.




FIG. 1: Bloch oscillations of $N = 5$ atoms in a parabolic lattice with parabolicity $\nu = 0.04J$: (a) mean momentum of non-interacting atoms; (b) mean momentum of interacting ($W = 0.2J$) atoms; (c) macroscopic coherence of the interacting atoms. Initial shift $l_0 = 8l^* = 80$.

The irreversible decay of BO according to Eq. (4) is a consequence of our approximation of the sum by an integral. Without this approximation, the decay of oscillations is followed by periodic revivals with a period $T_\nu = 2\pi\hbar/\nu$ [14]. One of these revivals is illustrated in the upper panel of Fig. 1, which shows the dynamics of the mean momentum of the non-interacting atoms in the parabolic lattice with parabolicity $\nu = 0.04J$. As initial state of the system we choose here the ground state of the atoms in a parabolic lattice with a slightly tighter confinement $\nu' = 4\nu$, which was then shifted by a distance $l_0 = 8l^* = 80$. Note that by changing ν' we change only the dephasing time [through the change of the wave packet width $\gamma = \gamma(\nu')$], while the revival time is defined exclusively by the parameter ν .

4. Next we address the effect of atom-atom interactions. In the case of quasi one-dimensional parabolic lattices, Bloch and dipole oscillations of the interacting atoms were studied in a number of papers, using the mean-field approach [13, 15, 16, 17, 18, 19]. As known, the mean-field approach is justified in the limit of large occupation number $\bar{n} \rightarrow \infty$ and vanishing microscopic interaction constant $W \rightarrow 0$, and leads (in the simplest case) to the discrete nonlinear Schrödinger equation,

$$i\hbar\dot{a}_l = \frac{\nu}{2}l^2a_l - \frac{J}{2}(a_{l+1} + a_{l-1}) + g|a_l|^2a_l, \quad (6)$$

where $g = WN$ is the macroscopic interaction constant. In our present work, we focus on the case of truly one-dimensional lattices, where the mean occupation number $\bar{n} \sim 1$. Clearly, the mean field approach is not applicable here and one has to treat the system microscopically by

using, for example, the Bose-Hubbard model,

$$H = \sum_l \frac{\nu}{2}l^2\hat{n}_l - \frac{J}{2} \left(\sum_l \hat{a}_{l+1}^\dagger \hat{a}_l + h.c. \right) + \frac{W}{2} \sum_l \hat{n}_l(\hat{n}_l - 1). \quad (7)$$

The main question we address below is the effect of atom-atom interactions on the Bloch dynamics depicted in the upper panel of Fig. 1.

First we shall discuss the initial conditions in some more detail. Throughout the paper we shall consider the ground many-body state of the atoms in a parabolic lattice as the initial wave packet (which is shifted then by the distance l_0). Clearly, along with the ratio J/ν this state is also defined by the ratio of the interaction constant to the hopping matrix element. Namely, it is essentially given by the symmetrized product of the single-particle atomic state for $W < J$, while it may resemble the Mott-insulator state for $W \gg J$ [12]. In what follows we restrict ourselves by a relatively weak interaction. Then the ground state of the system can be well approximated by the many-body wave function

$$|\tilde{\Psi}_0\rangle = \sum_{\mathbf{n}} c_{\mathbf{n}} |\mathbf{n}\rangle, \quad c_{\mathbf{n}} = \sqrt{N!} \prod_l \frac{a_l^{n_l}}{\sqrt{n_l!}}, \quad (8)$$

where $|\mathbf{n}\rangle = |\dots, n_{-1}, n_0, n_1, \dots\rangle$ is the Fock basis and the a_l satisfy the stationary nonlinear Schrödinger equation

$$\frac{\nu}{2}l^2a_l - \frac{J}{2}(a_{l+1} + a_{l-1}) + g|a_l|^2a_l = E_0a_l. \quad (9)$$

For example, for $N = 5$, $\nu = 0.04J$ and $W = 0.2J$, the overlap of the state (8) with the exact ground state $|\Psi_0\rangle$ is $|\langle\Psi_0|\tilde{\Psi}_0\rangle|^2 = 0.97$. We note that the state (8) is completely coherent and is analogous to the super-fluid state in a homogeneous lattice. We shall characterize the macroscopic coherence of the given many-body state $|\Psi\rangle$ by the maximal eigenvalue λ of the single-particle density matrix

$$\rho_{l,m} = N^{-1} \langle\Psi|\hat{a}_l^\dagger \hat{a}_m|\Psi\rangle. \quad (10)$$

Then the macroscopic coherence of the state (8) is $\lambda = 1$.

We proceed with the dynamics. The middle panel in Fig. 1 shows the mean momentum of $N = 5$ interacting atoms ($W = 0.2J$). In comparison with the noninteracting case (upper panel), a qualitative change is noticed. This change can be understood by analyzing the macroscopic coherence of the system, shown in the lower panel. It is seen that the macroscopic coherence oscillates with some characteristic period T_W . In the case of a homogeneously tilted lattice these oscillations were studied in Ref. [7]. The origin of the oscillations was shown to be the Stark localization of the single-particle wave functions which, together with the discreteness of the atom number, leads to the following expression for the macroscopic coherence,

$$\lambda = \exp(-2\bar{n}[1 - \cos(Wt/\hbar)]) . \quad (11)$$




FIG. 2: Dipole oscillations of $N = 4$ atoms in the parabolic lattice with parabolicity $\nu = 0.04J$: (a) mean momentum of non-interacting atoms; (b) mean momentum of interacting ($W = 0.2J$) atoms; (c) macroscopic coherence of interacting atoms. Initial shift $l_0 = l^*/2 = 5$.

In Eq. (11) $\bar{n}$ is the mean number of atoms per lattice site [20] and the limit $Fd \gg J$ is implicitly assumed. Since for the considered local static force $Fd = \nu l_0 = 3.2J$ Stark localization is not complete, the oscillations of the macroscopic coherence decay in time. Nevertheless, if this irreversible decay of coherence is slow on the time scale of the dephasing time, one can observe the revival of BO of the interacting atoms – an effect which attracts much attention because it provides an independent and accurate method for measuring the microscopic interaction constant W .

5. Let us now turn to the case $l_0 < l^*$. Here we meet dipole oscillations of a BEC with a characteristic frequency given by the frequency of small pendulum oscillations $\omega_0 = (\nu J)^{1/2}/\hbar$. (We recall in passing that the frequency of BO was given by $\omega_B = \nu l_0/\hbar \approx 2\omega_0 l_0/l^*$, $l_0 \gg l^*$.) For vanishing atom-atom interactions these dipole oscillations are shown in the upper panel of Fig. 2, where $l_0 = l^*/2 = 5$, and the other parameters are the same as in Fig. 1. The dephasing time τ_γ is again given by Eq. (5) but with the parameter ν substituted by the nonlinearity parameter $\tilde{\nu} = \nu/8$ [21]. (The latter parameter also defines the revival time.) The middle and lower panels in Fig. 2 refer to interacting atoms. An exponential decay of the macroscopic coherence is noticed. The other point to which we want to draw the attention of the reader is that a moderate interaction stabilizes the dipole oscillations against dephasing. Within the mean-

field approach (which reduces the Bose-Hubbard model to the discrete nonlinear Schrödinger equation), this phenomenon is discussed in Ref. [13].

6. In conclusion, we have shown that the dynamics of cold atoms in parabolic lattices is governed by the relation between two characteristic times – the dephasing time τ_γ and the decoherence time τ_W .

The dephasing time is inversely proportional to the width γ of the initial wave packet and the nonlinearity $\tilde{\nu}$, which, in turn, is defined by the initial shift l_0 of the wave packet relative to the separatrix $l^* = 2(J/\nu)^{1/2}$. Namely, $\tilde{\nu} = \nu/8$ for $l_0 \ll l^*$, and $\tilde{\nu} = \nu$ for $l_0 \gg l^*$. It is interesting to estimate the dephasing time in a typical laboratory experiment. Taking, as an example, the recent experiment [10] with rubidium atoms in an array of axially modulated quantum tubes, we have $\nu = 0.0014E_R$ and $J = 0.38E_R$ for the modulation amplitude (depth of the optical lattice) of one recoil energy. This gives a separatrix position $l^* = 33$, and a period $T_0 = 12.1$ ms of small dipole oscillations. Assuming a dilute gas (which, in fact, is not the case realized in the cited experiment) the width of the initial wave packet is $\gamma \approx (J/4\nu)^{1/4} = \sqrt{l^*}/2 \approx 3$ and, hence, the dephasing time $\tau_\gamma = 85$ ms for dipole oscillations, and $\tau_\gamma = 10.6$ ms for BO. Note that these are upper estimates for the dephasing times, and for the initial shift l_0 closer to the separatrix the dephasing times are essentially smaller. It is also worth noting that there is a maximal shift l_0 above which the single band approximation (used throughout the paper) is not valid. The crucial parameter here is the energy gap between the Bloch bands ($\Delta = 0.5E_R$ for the specified parameters). The analysis of BO in a parabolic lattice beyond the single-band approximation will be subject of a separate paper.

The decoherence time τ_W is defined by the characteristic density of the atomic gas $\bar{n}$ and by the value of the microscopic interaction constant W . The latter, in turn, is defined by the s -wave scattering length and by the degree of confinement of the atoms in the wells of the optical potential. In particular, in the experiment [10], the quantum tubes were created by two crossing quasi 1D optical lattices with an amplitude $V = 30E_R$. For the axial modulation with $V = E_R$ this gives $W = 0.73E_R$. For this relatively high value of the interaction constant few atoms per one tube are enough to destroy the dipole/Bloch oscillations on a very short time scale. This qualitatively explains the results of the experiment [10], where the number of atoms per one quantum tube was around 20. To observe the effects discussed in this paper one has to decrease either the atomic density or the interaction constant (e.g. by use of a Fishbach resonance), as compared to those of Ref. [10].

[1] C. Zener, Proc. R. Soc. Lond. **A145**, 523 (1934).

[2] M. Ben Dahan, E. Peik, J. Reichel, Y. Castin, and C. Sa-

lomon, Phys. Rev. Lett. **76**, 4508 (1996).

[3] For the introductory review see A.R.Kolovsky and

- H.J.Korsch, International J. of Mod. Physics **18**, 1235 (2004).
- [4] O. Morsch, J. H. Müller, M. Cristani, D. Ciampini, and E. Arimondo, Phys. Rev. Lett. **87**, 140402 (2001).
 - [5] R. G. Scott, A. M. Martin, S. Bujkiewicz, T. M. Fromhold, N. Malossi, O. Morsch, M. Cristiani, and E. Arimondo, Phys. Rev. A **69**, 033605 (2004).
 - [6] M. Modugno, E. de Mirandes, F. Ferlaino, H. Ott, G. Roati, and M. Inguscio, Fortschr. Phys. **52**, 1173 (2004).
 - [7] A. R. Kolovsky, Phys. Rev. Lett. **90**, 213002 (2003).
 - [8] A. Buchleitner and A. R. Kolovsky, Phys. Rev. Lett. **91**, 253002 (2003).
 - [9] A. R. Kolovsky and A. Buchleitner, Phys. Rev. E **68**, 056213 (2003).
 - [10] C. D. Fertig, K. M. O'Hara, J. H. Huckans, S. L. Rolston, W. D. Phillips, and J. V. Porto, Phys. Rev. Lett. **94**, 120403 (2005).
 - [11] F. S. Cataliotti, L. Fallani, F. Ferlaino, C. Fort, P. Madaloni, and M. Inguscio, New J. of Phys. **5**, 71.1 (2003).
 - [12] A. M. Rey, G. Pupillo, C. W. Clark, and C. J. Williams, e-print: cond-mat/0503477.
 - [13] J. Brand and A. R. Kolovsky, e-print: cond-mat/0412549.
 - [14] Indeed, at times t multiple to T_ν , the phases $\nu mt/\hbar$ in the sum (3) are multiple of 2π and, hence, the momentum takes its initial value. (More rigourosly, the revivals of ocsillations follow from the quadratic dependence for the eigenenergies of the quantum pendulum in the asymptotic region $l \gg l^*$.)
 - [15] M. Krämer, L. Pitaevskii, and S. Stringari, Phys. Rev. Lett. **88**, 180404 (2002).
 - [16] M. L. Chiofalo and M. P. Tosi, J. Phys. B: At. Mol. Opt. Phys. **34**, 4551 (2001).
 - [17] A. Smerzi, A. Trombettoni, P. G. Kevrekidis, and A. R. Bishop, Phys. Rev. Lett. **89**, 170402 (2002).
 - [18] S. K. Adhikari, Eur. Phys. J. D **25**, 161 (2003).
 - [19] F. Nesi and M. Modugno, J. Phys. B: At. Mol. Opt. Phys. **37**, S101 (2004).
 - [20] In the considered case of a finite wave packet it looks reasonable to associate $\bar{n}$ with the number of atoms N devided by the wave packet width γ .
 - [21] A. J. Lichtenberg and M. A. Leibermann, *Regular and chaotic dynamics* (Springer, Berlin, 1983).